



BILL & MELINDA GATES INSTITUTE for POPULATION and REPRODUCTIVE HEALTH

Niger (Niamey) Snapshot of Indicators Phase 1 (2021) Cross-sectional Survey











Background Characteristics of Respondents

Distribution of women age 15 to 49 by background characteristics.

Background	Percentage	Weighted	Unweighted	
Characteristics	of Sample	Number	Number	
Total				
All	100.0	274	1355	
Age Group				
15-19	23.9	66	329	
20-24	19.8	54	267	
25-29	16.2	44	224	
30-34	13.2	36	179	
35-39	12.7	35	166	
40-44	8.4	23	116	
45-49	5.7	16	74	
Marital Status				
Married, or in union	59.9	164	789	
Unmarried, sexually active	0.9	2	13	
Parity				
0-1 children	11.8	32	161	
2-3 children	24.1	66	324	
4+ children	25.6	70	335	
Residence				
Urban	94.4	259	1264	
Rural	5.6	15	91	
Education				
Never	27.2	74	354	
Primary	20.1	55	276	
Secondary	38.9	107	534	
Tertiary	13.8	38	190	
Wealth Quintile				
Lowest	0.3	1	4	
Middle	3.1	8	41	
Highest	96.6	265	1310	
•				



Current Use of Contraception

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics.

	Any N	lethod	Modern	Method ¹	Traditiona	Traditional Method		
Background Characteristics	All Women	Married ² Women	All Women	Married ² Women	All Women	Married ² Women		
Total								
All	20.0	31.6	19.1	30.2	0.8	1.4		
Age Group								
15-19	4.4	24.4	4.1	22.5	0.3	1.9		
20-24	17.1	33.2	16.6	32.1	0.5	1.1		
25-29	36.0	43.9	34.3	41.7	1.7	2.2		
30-34	36.1	40.2	34.4	38.3	1.7	2.0		
35-39	23.3	25.7	22.1	24.2	1.2	1.4		
40-44	15.4	16.0	15.4	16.0	0.0	0.0		
45-49	11.1	14.7	11.1	14.7	0.0	0.0		
Marital Status								
Married, or in union	31.6	na	30.2	na	1.4	na		
Unmarried, sexually active		na		na		na		
Parity								
0-1 children	24.7	30.1	22.3	27.0	2.4	3.0		
2-3 children	39.3	41.8	37.0	39.3	2.3	2.6		
4+ children	28.4	29.6	28.4	29.6	0.0	0.0		
Residence								
Urban	19.9	31.9	19.0	30.4	0.9	1.5		
Rural	20.8	27.4	20.8	27.4	0.0	0.0		
Education								
Never	20.5	24.0	19.6	23.0	0.9	1.1		
Primary	23.7	31.0	23.7	31.0	0.0	0.0		
Secondary	16.1	35.8	15.8	34.9	0.4	0.9		
Tertiary	24.3	45.7	21.0	39.5	3.3	6.2		
Wealth Quintile								
Lowest								
Middle	14.9	23.7	14.9	23.7	0.0	0.0		
Highest	20.2	32.0	19.3	30.6	0.9	1.5		

¹ Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

² Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Contraception Method Mix

Percent distribution of current contraceptive users age 15 to 49, by method type, marital status, and background characteristics.

Background Characteristics

All Users (N = 274)

Married Users¹ (N = 260)

Characteristics						, (-,			_					1		,		
9	Sterilization	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl	Steriliztn	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtn
Total																				
All	0.7	20.3	2.7	4.4	10.6	53.4	0.3	2.5	1.0	4.2	0.7	19.1	2.9	4.6	10.8	55.0	0.3	1.1	1.1	4.4
Age Group																				
15-19																				
20-24	0.0	26.7	0.0	9.7	9.7	49.5	0.0	1.3	0.0	3.2	0.0	27.8	0.0	10.1	10.1	48.7	0.0	0.0	0.0	3.3
25-29	0.0	23.9	3.5	0.8	11.1	52.9	0.9	2.2	0.0	4.7	0.0	22.1	3.7	0.9	11.8	54.4	1.0	1.0	0.0	5.0
30-34	0.0	17.3	0.9	2.3	11.2	61.8	0.0	0.0	1.6	4.8	0.0	16.3	0.9	2.4	11.4	62.6	0.0	0.0	1.6	4.9
35-39	0.0	14.6	3.3	8.0	13.9	52.9	0.0	0.0	2.2	5.2	0.0	11.5	3.5	8.5	14.7	53.9	0.0	0.0	2.4	5.5
40-44																				
45-49																				
Marital Status																				
Married, or in union	0.7	19.1	2.9	4.6	10.8	55.0	0.3	1.1	1.1	4.4	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually ac	tive										na	na	na	na	na	na	na	na	na	na
Parity																				
0-1 children	0.0	25.3	0.0	3.9	2.2	48.8	0.0	7.4	2.6	9.9	0.0	25.9	0.0	3.9	2.3	50.0	0.0	5.2	2.6	10.1
2-3 children	0.0	22.1	3.6	4.0	7.9	55.2	0.0	1.4	0.0	5.8	0.0	20.7	3.8	4.2	8.4	56.1	0.0	0.7	0.0	6.1
4+ children	1.9	17.0	2.7	5.3	17.0	54.4	0.0	0.0	1.8	0.0	1.9	14.5	2.8	5.5	17.7	55.7	0.0	0.0	1.9	0.0
Residence																				
Urban	0.7	20.6	2.6	4.6	8.3	54.7	0.3	2.6	1.1	4.4	0.8	19.3	2.7	4.9	8.4	56.5	0.3	1.2	1.2	4.7
Rural																				
Education																				
Never	1.4	16.4	2.4	5.1	7.4	61.3	0.0	1.8	0.0	4.2	1.5	16.3	2.6	5.5	6.6	63.1	0.0	0.0	0.0	4.4
Primary	0.0	15.9	3.1	7.2	23.8	47.2	0.0	1.4	1.4	0.0	0.0	15.5	3.3	7.6	25.4	46.7	0.0	0.0	1.5	0.0
Secondary	0.9	26.3	2.5	3.9	6.2	52.6	0.0	4.2	1.2	2.3	0.9	22.6	2.7	4.2	6.6	56.6	0.0	2.5	1.3	2.5
Tertiary	0.0	22.0	2.9	0.0	5.4	50.5	1.7	2.0	1.9	13.7	0.0	22.0	2.9	0.0	5.4	50.5	1.7	2.0	1.9	13.7
Wealth Quintile	e																			
Lowest	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na
Middle																				
Highest	0.7	20.2	2.1	4.5	10.2	54.3	0.3	2.5	1.1	4.3	0.7	18.8	2.2	4.7	10.4	56.0	0.3	1.2	1.1	4.5
											1									

¹ Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Modern contraceptive methods

Percentage of women ages 15 to 49 using short-acting and long-acting contraceptive methods by background characteristics.

Background Characteristics

Characteristics	Short Acti	ing Methods ¹	Long Actii	ng Methods ²
	All Women	Married Women ³	All Women	Married Women ³
	(N = 1355)	(N = 789)	(N = 1355)	(N = 789)
Total				
All	14.4	23.0	4.7	7.1
Age Group				
15-19	3.5	18.5	0.6	4.0
20-24	12.0	22.9	4.6	9.2
25-29	24.5	30.4	9.8	11.3
30-34	27.8	31.3	6.6	7.0
35-39	18.0	20.4	4.2	3.9
40-44	9.3	10.2	6.2	5.8
45-49	8.2	10.8	2.9	3.9
Marital Status				
Married, or in union	23.0	na	7.1	na
Unmarried, sexually active		na		na
Parity				
0-1 children	16.0	19.3	6.2	7.8
2-3 children	26.9	29.0	10.1	10.3
4+ children	22.3	23.9	6.1	5.7
Residence				
Urban	14.3	23.1	4.8	7.3
Rural	16.4	21.6	4.4	5.8
Education				
Never	15.5	18.1	4.1	4.9
Primary	19.2	25.1	4.5	5.8
Secondary	11.0	25.5	4.8	9.4
Tertiary	14.9	28.1	6.1	11.4
Wealth Quintile				
Lowest				
Middle	6.3	10.0	8.6	13.7
Highest	14.7	23.6	4.6	7.0

¹ Short-acting methods include injectables, pills, emergency contraception, condom (male and female), standard days/cycle beads, and LAM.

² Long-acting methods include IUD, implants, and sterilization (male and female).

³ Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Unmet Need for Family Planning

Percent of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics.

Background Characteristics

Background Characteristics	All V	Vomen (N = 13	55)	Married Women ¹ (N = 789)			
	For Spacing	For Limiting	Total	For Spacing	For Limiting	Total	
Total							
All	9.5	1.3	10.8	15.3	2.2	17.5	
Age Group							
15-19	3.2	0.0	3.2	21.0	0.0	21.0	
20-24	7.9	0.0	7.9	14.7	0.0	14.7	
25-29	11.9	0.0	11.9	15.1	0.0	15.1	
30-34	14.3	0.7	15.0	16.1	0.8	16.9	
35-39	14.8	2.5	17.3	17.3	2.9	20.2	
40-44	12.0	6.7	18.7	11.9	8.1	20.0	
45-49	8.0	5.7	13.6	10.5	7.5	18.0	
Marital Status							
Married, or in union	15.3	2.2	17.5	na	na	na	
Unmarried, sexually active				na	na	na	
Parity							
0-1 children	13.7	0.8	14.5	16.5	1.0	17.5	
2-3 children	13.3	1.1	14.4	14.8	1.2	16.0	
4+ children	17.1	3.7	20.8	17.8	4.0	21.8	
Residence							
Urban	9.7	1.4	11.0	15.8	2.3	18.1	
Rural	6.9	0.0	6.9	9.0	0.0	9.0	
Education							
Never	13.7	2.4	16.0	16.7	3.0	19.6	
Primary	13.1	1.7	14.8	16.7	2.4	19.1	
Secondary	7.2	0.3	7.5	16.8	0.8	17.6	
Tertiary	2.7	1.3	3.9	5.0	2.4	7.4	
Wealth Quintile							
Lowest							
Middle	2.1	0.0	2.1	3.3	0.0	3.3	
Highest	9.7	1.3	11.0	15.6	2.3	17.9	

¹ Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i>Italics indicate estimate was generated from 25 to 49 unweighted observations



Need and Demand for Family Planning

Percent of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics.

Married Women² (N=789)

Background Characteristics

Cital acteristics		All W	omen (r	N=1322)			iviarried	women	(N=789)	
	Any Method	Modern ¹ Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern	Any Method	Modern ¹ Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern
Total										
All	20.0	19.1	10.8	30.7	62.2	31.6	30.2	17.5	49.0	61.5
Age Group										
15-19	4.4	4.1	3.2	7.6	54.1	24.4	22.5	21.0	45.4	
20-24	17.1	16.6	7.9	25.0	66.3	33.2	32.1	14.7	47.9	67.1
25-29	36.0	34.3	11.9	47.9	71.6	43.9	41.7	15.1	59.0	70.7
30-34	36.1	34.4	15.0	51.1	67.3	40.2	38.3	16.9	57.1	67.0
35-39	23.3	22.1	17.3	40.6	54.5	25.7	24.2	20.2	45.9	52.9
40-44	15.4	15.4	18.7	34.1	45.1	16.0	16.0	20.0	36.0	44.4
45-49	11.1	11.1	13.6	24.8		14.7	14.7	18.0	32.7	
Marital Status										
Married, or in union	31.6	30.2	17.5	49.0	61.5	na	na	na	na	na
Unmarried, sexually active						na	na	na	na	na
Parity										
0-1 children	24.7	22.3	14.5	39.2	56.8	30.1	27.0	17.5	47.6	56.8
2-3 children	39.3	37.0	14.4	53.7	68.9	41.8	39.3	16.0	57.8	67.9
4+ children	28.4	28.4	20.8	49.2	57.7	29.6	29.6	21.8	51.4	57.6
Residence										
Urban	19.9	19.0	11.0	30.9	61.5	31.9	30.4	18.1	50.0	60.8
Rural	20.8	20.8	6.9	27.7	75.2	27.4	27.4	9.0	36.4	75.2
Education										
Never	20.5	19.6	16.0	36.5	53.8	24.0	23.0	19.6	43.7	52.6
Primary	23.7	23.7	14.8	38.5	61.5	31.0	31.0	19.1	50.1	61.8
Secondary	16.1	15.8	7.5	23.6	66.7	35.8	34.9	17.6	53.4	65.4
Tertiary	24.3	21.0	3.9	28.2	74.3	45.7	39.5	7.4	53.1	74.3
Wealth Quintile										
Lowest										
Middle	14.9	14.9	2.1	17.0		23.7	23.7	3.3	27.0	
Highest	20.2	19.3	11.0	31.2	61.9	32.0	30.6	17.9	49.9	61.3

All Women (N=1355)

¹ Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

² Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Pregnancy Intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics.

Background	All Wom	en with 1 or N	More Births	Married Wo	men with 1 o	r More Births
Characteristics		(N=598)			(N=561)	
	Wanted Then	Wanted Later	Not At All	Wanted Then	Wanted Later	Not At All
Total						
All	79.3	19.3	1.4	79.8	18.8	1.3
Age Group						
15-19	75.5	24.5	0.0	74.4	25.6	0.0
20-24	70.6	29.4	0.0	71.6	28.4	0.0
25-29	85.2	14.8	0.0	86.4	13.6	0.0
30-34	83.6	14.0	2.4	83.4	14.2	2.5
35-39	72.1	27.4	0.5	72.5	27.0	0.6
40-44	87.4	4.9	7.7	88.4	5.3	6.3
45-49						
Marital Status						
Married, or in union	79.8	18.8	1.3	na	na	na
Unmarried, sexually active				na	na	na
Parity						
0-1 children	83.6	16.4	0.0	84.9	15.1	0.0
2-3 children	79.7	19.6	0.7	80.3	19.0	0.7
4+ children	76.9	20.3	2.8	77.2	20.3	2.5
Residence						
Urban	79.1	19.4	1.5	79.7	18.9	1.4
Rural	81.7	18.3	0.0	81.7	18.3	0.0
Education						
Never	79.8	19.3	0.9	79.7	19.8	0.5
Primary	80.0	19.1	0.8	80.1	19.0	0.9
Secondary	77.5	20.9	1.6	80.0	18.3	1.7
Tertiary	80.1	16.1	3.7	79.2	16.9	3.9
Wealth Quintile						
Lowest						
Middle						
Highest	79.3	19.3	1.5	79.9	18.7	1.4
	, 5.5	13.3	1.5	, , , ,	10.7	1.7

¹ Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i>Italics indicate estimate was generated from 25 to 49 unweighted observations



Contraceptive Choice

Percent distribution of women age 15 to 49 who are currently using modern contraception reporting on who decided on the contraceptive method, by background characteristics.

Background Characteristics	Self	Jointly with P Partner/Provider	rovider/Partner Alone	(N = 260)
Total				
All	65.7	28.4	5.9	
Age Group				
15-19				
20-24	74.3	24.3	1.4	
25-29	64.9	29.6	5.6	
30-34	55.8	39.2	5.0	
35-39	68.3	22.5	9.2	
40-44				
45-49				
Marital Status				
Married, or in union	65.6	29.0	5.4	
Unmarried, sexually active				
Parity				
0-1 children	64.7	32.9	2.3	
2-3 children	70.2	27.1	2.7	
4+ children	60.9	29.8	9.3	
Residence				
Urban	64.5	30.3	5.2	
Rural				
Education				
Never	64.4	28.0	7.6	
Primary	71.3	26.5	2.2	
Secondary	63.5	28.2	8.3	
Tertiary	63.7	32.9	3.4	
Wealth Quintile				
Lowest	na	na	na	
Middle				
Highest	64.8	29.1	6.0	
5				

¹ Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i>Italics indicate estimate was generated from 25 to 49 unweighted observations



Method Information Index Components

Percentage of current modern contraceptive ¹ users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics.

		about other ² thods		about side 2 ects	Told what to do side e	if experienced 4 ffects		t switching 2 s in future
Background Characteristics	All	Married ³	All	Married ³	All	Married ³	All	Married ³
	Users	Users	Users	Users	Users	Users	Users	Users
Total	(260)	(246)	(260)	(246)	(145)	(140)	(260)	(246)
All	66.1	68.0	55.5	56.5	85.5	85.0	64.0	65.6
Age Group								
15-19								
20-24	73.1	73.4	53.6	55.9	85.7	85.7	61.1	63.8
25-29	63.8	68.0	60.9	61.5	81.4	80.3	64.0	66.0
30-34	70.7	70.4	50.9	51.5	95.0	95.0	68.2	67.8
35-39	71.2	71.7	60.2	60.1			75.2	76.0
40-44								
45-49								
Marital Status								
Married, or in union	68.0	na	56.5	na	85.0	na	65.6	na
Unmarried, sexually active		na		na	na	na		na
Parity								
0-1 children	85.9	88.3	59.3	61.0			76.0	78.1
2-3 children	62.4	64.2	53.3	54.3	86.1	85.6	58.1	59.4
4+ children	66.0	65.5	58.6	57.8	84.7	83.8	69.3	69.0
Residence								
Urban	65.3	67.2	54.7	55.8	86.9	86.3	62.7	64.3
Rural			J-1.7					
Education								
Never	62.7	63.6	61.0	64.8	81.0	81.0	64.0	66.9
Primary	66.4	69.3	50.7	52.6	80.3	79.8	67.4	70.4
•	69.4	72.1		52.6	90.6			
Secondary Tertiary	65.4	65.4	54.0 56.2	56.2	90.0	89.5 	61.6 <i>63.7</i>	61.7 <i>63.7</i>
,	03.4	03.4	30.2	30.2			03.7	03.7
Wealth Quintile								
Lowest	na	na	na	na	na	na	na	na
Middle				 FC F		 0F 3	 C2 F	 CE 4
Highest	66.0	67.9	55.4	56.5	85.8	85.2	63.5	65.1

¹ Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

- 2 Asked of all current/recent modern contraceptive users
- 3 Married includes married women and those who are living with a man
- 4 Asked of all current/recent modern contraceptive users who were informed about side effects
- na Not applicable
- -- Less than 25 unweighted observations
- <i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Quality of Care

Percentage of current modern¹contraceptive users who would return, would refer a friend/relative, or would return and refer a friend/relative to provider, by background characteristics.

Background Characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Total All	(N = 256) 86.6	(N = 256) 84.3	(N = 256) 78.1
Age Group			
15-19			
20-24	91.9	79.6	78.3
25-29	86.7	89.5	83.6
30-34	90.7	93.1	83.8
35-39	82.7	81.4	72.6
40-44			
45-49			
Marital Status			
Married, or in union	87.2	84.9	79.2
Unmarried, sexually active			
Parity			
0-1 children	84.9	87.2	76.8
2-3 children	85.3	84.5	78.1
4+ children	89.0	84.6	80.0
	33.3	0.110	55.5
Residence			
Urban	87.1	84.3	78.1
Rural			
Education			
Never	90.2	84.0	80.2
Primary	76.7	83.8	72.1
Secondary	90.5	88.7	84.9
Tertiary	87.5	76.6	69.7
Wealth Quintile			
Lowest	na	na	na
Middle			
Highest	86.6	84.3	77.9

¹ Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Knowledge of Family Planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics.

Background Characteristics	Visited by heal discussed fai	mily planning	Visited facility discussed far	mily planning	Either visited by or visited f discussed far	acility and mily planning
	All Women	Married ¹ Women	All Women	Married ¹ Women	All Women	Married ¹ Women
Total	(1355)	(789)	(1355)	(789)	(1355)	(789)
All	3.1	3.9	12.3	19.5	14.4	22.2
Age Group						
15-19	1.7	5.3	4.9	29.9	6.3	33.4
20-24	2.4	3.4	10.5	18.7	11.7	21.0
25-29	3.7	4.2	19.4	24.6	21.9	27.7
30-34	2.6	2.3	19.4	21.8	21.6	23.7
35-39	7.0	5.9	18.8	20.3	23.3	23.4
40-44	3.2	3.1	7.9	8.7	10.5	11.8
45-49	2.6	3.5	5.7	7.6	6.8	9.0
Marital Status						
Married, or in union	3.9	na	19.5	na	22.2	na
Unmarried, sexually active		na		na		na
Parity						
0-1 children	4.0	3.1	19.2	23.4	22.1	25.0
2-3 children	3.5	3.9	19.9	21.9	22.2	24.5
4+ children	4.9	4.8	17.9	19.2	21.3	22.6
Residence						
Urban	2.1	2.5	11.5	18.5	12.6	19.8
Rural	20.5	22.5	26.5	33.5	44.8	53.0
Education						
Never	5.9	5.9	15.2	17.9	19.4	22.0
Primary	2.7	3.4	17.8	23.7	19.1	25.5
Secondary	1.4	2.1	8.6	19.9	9.8	21.6
Tertiary	2.9	2.7	9.5	15.3	11.2	17.5
Wealth Quintile						
Lowest						
Middle	24.8	26.6	11.8	15.9	30.6	35.9
Highest	2.4	3.0	12.3	19.6	13.8	21.6
	 .	0.0		20.0	20.0	

¹ Married includes married women and those who are living with a man

na Not applicable

⁻⁻ Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations



Response Rates

Response rates of households and individuals, by residence.

Result	Urban	Rural	Total
Household Interviews			
Households Selected	1085	70	1155
Households Occupied	1060	70	1130
Households Interviewed	1035	70	1105
Household Response Rate ¹	97.64%	100.00%	97.79%
Interviews with Women			
Number of Eligible Women ²	1298	95	1393
Number of Eligible Women Interviewed	1264	91	1355
Eligible Woman Response Rate ³	97.38%	95.79%	97.27%

¹ Household response rate = households interviewed/households selected

² Eligible women are women ages 15-49 identified in completed household interviews

³ Eligible response rate = eligible women interviewed/eligible women



Sample Error Estimates

	All Women					Married Women ¹			
Variable	Mean	Std Error	CI Lower	CI Upper	Mean	Std Error	CI Lower	CI Upper	
Currently using a modern method	0.191	0.012	0.168	0.217	0.302	0.020	0.262	0.345	
Currently using a traditional method	0.008	0.003	0.004	0.016	0.014	0.005	0.007	0.028	
Currently using any contraceptive method	0.200	0.013	0.174	0.227	0.316	0.022	0.272	0.363	
Currently using DMPA-IM	0.044	0.013	0.023	0.080	0.046	0.014	0.024	0.085	
Currently using DMPA-SC	0.106	0.023	0.067	0.163	0.108	0.024	0.068	0.168	
Currently using condoms	0.025	0.010	0.011	0.054	0.011	0.006	0.004	0.033	
Currenlty using implants	0.203	0.022	0.162	0.253	0.191	0.025	0.146	0.246	
Chose method by self or jointly in past 12 months	0.941	0.022	0.876	0.973	0.946	0.019	0.890	0.974	
Informed by provider about other methods	0.661	0.041	0.574	0.739	0.680	0.042	0.588	0.759	
Informed by provider about side effects	0.555	0.047	0.459	0.647	0.565	0.047	0.469	0.657	
Satisfied with provider: Would return and refer friend/relative to provider	0.781	0.045	0.675	0.860	0.792	0.045	0.687	0.869	
Visited by health worker who talked about family planning in past 12 months	0.031	0.011	0.015	0.064	0.039	0.016	0.017	0.085	

¹ Married includes married women and those who are living with a man